


MISR Retrieval Examples: Red Sea, 25 March & 29 June 2001



70°-forward images

558-nm aerosol optical depth


The MISR Program for Aerosol Product Intensive Validation

Validate MISR Aerosol Retrieval Algorithms

Field Campaigns provide some of our best opportunities to

- Test Our Retrieval Assumptions
 with the help of the aircraft- and surface-based aerosol observations
- Refine Our Algorithms Accordingly

We Must Address Key Algorithm Assumptions



We Must Identify Specific Sites where data offer critical tests

The MISR Program for Aerosol Product Intensive Validation (Cont'd)

Constrain Regional Transports and Radiative Forcing

Once we have a Validated Algorithm, MISR will provide over **Entire Regions**, on a **Regular Basis**

- Aerosol Column Optical Depth
- A Constraint on Aerosol Type

This includes

- Re-running MISR data from CLAMS and other Campaigns
- Comparisons with **Transport Models**
- Global Radiative Forcing Calculations

MISR Team Goals for Participation in the CLAMS Summer 2001 IFC

1. MISR Retrieval Validation -- to test our multi-angle aerosol retrieval approaches over dark water, at least using AirMISR observations, but we hope also with MISR observations.

To meet this goal, we need as close to a "**column closure**" experiment as possible. The aim is to characterize at least one column in as much detail as possible -- aerosol properties, vertical distribution, radiation field, boundary conditions -- coincident with AirMISR (and possibly MISR) overflight, under cloud-free conditions.

We anticipate testing our sensitivity to maritime aerosol air masses, possibly clean continental or industrially polluted aerosol air masses, and thin cirrus at COVE.


The collection of aircraft and surface instruments involved makes this a unique opportunity to meet one of our key validation goals.

2. Scene Variability -- to quantitatively assess the contribution of sub-MISR-pixel scene variability to aerosol measurement uncertainty over dark water.


To meet this goal, we need a "Volume Closure" experiment, with both MISR and AirMISR observing the region under cloud-free conditions. At the same time, the aggregate of surface and aircraft platforms have to do the best job they can characterizing the spatial variability of the surface BRDF and atmospheric gas and aerosols, on one-to-ten-kilometer scales.


Since there is **only about a 1 in 3 chance of cloud-free conditions** for a MISR overpass during the Summer 2001 IFC, it is less likely we will meet this goal. We may reassess our options for meeting this goal next year.

MISR paths over Chesopeake, Local Mode Site #040 (36.900, -75.710)



MISR Local Mode Regions for CLAMS






Orbit	X-Track	Of Time	Of Date
§7708	163.1 km	18:09:50	30Mcy01
£7941	153.5 km	18:09:01	15Jun01
#8174	161.0 km	16:10:00	0130001
#8407	161.0 km	16:10:00	1730/01
#8640	161.0 km	16:10:00	02Aug01

Neorby LM Sites

179 USDA_MD

184 Wollops_Is

187 Woshingtn_DC



MISR coverage During the CLAMS Campaign July 10- August 1, 2001

Date	UTC	Target	Orbit	Path
July 10 2001	191_16:06	Chesapeake	8305	14
July 17 2001	198_16:12	Chesapeake	8407	15
July 19 2001	200_16:00	Chesapeake	8436	13
July 26 2001	207_16:06	Chesapeake	8538	14
Aug 2 2001	214_16:12	Chesapeake	8640	15
Aug 4 2001	216_16:00	Chesapeake	8669	13

...of which, Path 15 is less good than 13 and 14

MISR and AirMISR at CLAMS

	<u>AirMISR</u>	MISR
Platform	ER2 Aircraft	Terra Satellite
Elevation	20 km	704 km
Angular Coverage	{±70.5, ±60°, ±45.6°, ±26.1°, nadir}	same
Spectral Channels	{446, 558, 672, 867 nm}	same
Pixel Resolution	27.5 m	275 m
9-Angle Coverage	11 x 9 km Patch	360-km Swath
Observing Method	1 Pivoting Camera	9 Cameras

AirMISR acquires 9 images over the range of angles in a **12-min**, **147-km** ER-2 flight line.

MISR acquires 9 images of a 360-km swath over a period of 7 minutes, about once per week at COVE.

MISR **CLAMS** Validation Days

[ALL MISR Results are as yet Preliminary – **UNVALIDATED**]

July 10 -- Chesapeake Lighthouse, Orbit 8305, 16:02:32 UTC

Cloud-free day; CV-580 deployed 3 hrs. after MISR overflight.

MISR – Industrial Maritime (80% medium, 10% large, 10% BC), Tau = 0.272

July 12 -- Chesapeake Lighthouse, AirMISR only

Cloud-free run with AirMISR over Lighthouse, and simultaneously, CV-580

**<u>July 17</u> -- Chesapeake Lighthouse, Orbit 8407, 16:08:37 UTC

Golden Day – 5 instrumented aircraft, including ER-2 carrying AirMISR, stacked over Lighthouse at MISR pass MISR – Clean Maritime (70% medium; 15% large; 15% very large), Tau = 0.443

July 26 -- Chesapeake Lighthouse, Orbit 8538, 16:02:29 UTC

Cloudy day – MISR overflew with CV-580 in a "clear" patch east of Lighthouse MISR – Clean Maritime (70% medium; 15% large; 15% very large), Tau = 0.535


July 31 -- Chesapeake Lighthouse, AirMISR only

AirMISR flew star patterns over deep water buoy as other aircraft measured BRDF


**<u>Aug. 02</u> -- Chesapeake Lighthouse, Orbit 8640, 16:08:28 UTC

Golden Day – During MISR pass, AirMISR and CV-580 overflew Lighthouse


CLAMS: Orbit 8305, Block 61 July 10, 2001 MISR Level 1B2 ELLIPSOID RGB AN




CLAMS: Orbit 8407, Block 61 July 17, 2001 MISR Level 1B2 ELLIPSOID RGB AN




CLAMS: Orbit 8538, Block 61 July 26, 2001 MISR Level 1B2 ELLIPSOID RGB AN




CLAMS: Orbit 8640, Block 61 August 2, 2001 MISR Level 1B2 ELLIPSOID RGB AN



MISR Dark Water Calibration Using CLAMS AirMISR Data





MISR dark scene calibration exercise using coincident MISR and AirMISR data over ocean near the Chesapeake Lighthouse during the CLAMS field campaign. Such data were taken on a high-aerosol-optical-depth day (July 17, 2001, shown above), when mid-visible optical depths were on order 0.5, an on August 2, 2001, when the mid-visible aerosol optical depth was about 0.15. Excluding the cameras affected by sun glint (3, 4, 5, and in the NIR, 6 for the cases shown above), the MISR data was systematically brighter than AirMISR on both days. The discrepancies are between 5% and 8%. This was traced to ghosting in the MISR optics, which affects the brightness of dark targets (such as ocean) in scenes that also contain bright features (land, sea ice, clouds). The MISR Team is working to develop and implement corrections.

Correlative Ocean Surface Measurements Made During CLAMS


Date	MISR Time†	AirMISR Time†	CAR Time†	Site
7/10	16:02		18:12*	COVE
7/12		15:37	12:21	COVE
7/17	16:08	16:02	16:57	COVE
7/31		15:44-16:43	16:59	Near buoy
		(star)		44004
8/2	16:08	16:02	20:10**	COVE

^{*}CV-580 takeoff delayed 3 hours waiting for clearance from LaRC safety officer.

^{**}CAR door stuck closed during MISR overflight.

[†] Times are UTC.

CLAMS Field Validation Campaign



Chesapeake Lighthouse, VA, July 17, 2001